

Automated Reasoning - WS 2013

Lecture 4: Examples

Mădălina Erăscu and Tudor Jebelean
 Research Institute for Symbolic Computation,
 Johannes Kepler University, Linz, Austria
`{merascu,tjebeloa}@risc.jku.at`

October 30 & November 6, 2013

Example 1 (Classification) Transform the formulae F_1, F_2, F_3, F_4 , and $\neg G$ into a set of clauses, where

$$\begin{aligned}
 F_1 : \quad & \forall_{x,y} \exists_z P[x,y,z] \\
 & \forall_{x,y,z,u,v,w} (P[x,y,u] \wedge P[y,z,v] \wedge P[u,z,w] \Rightarrow P[x,v,w]) \\
 F_2 : \quad & \wedge \\
 & \forall_{x,y,z,u,v,w} (P[x,y,u] \wedge P[y,z,v] \wedge P[x,v,w] \Rightarrow P[u,z,w]) \\
 F_3 : \quad & \forall_x P[x,e,x] \wedge \forall_x P[e,x,x] \\
 F_4 : \quad & \forall_x P[x,i[x],e] \wedge \forall_x P[i[x],x,e] \\
 G : \quad & \left(\forall_x P[x,x,e] \right) \Rightarrow \left(\forall_{u,v,w} (P[u,v,w] \Rightarrow P[v,u,w]) \right)
 \end{aligned}$$

Solution. F_1, F_2, F_3, F_4 can almost immediately transformed into clauses. We have

$$\begin{aligned}
 & P[x,y,f[x,y]] \\
 & \neg P[x,y,u] \vee \neg P[y,z,v] \vee \neg P[u,z,w] \vee P[x,v,w] \\
 & \neg P[x,y,u] \vee \neg P[y,z,v] \vee \neg P[x,v,w] \vee P[u,z,w] \\
 & P[x,e,x] \\
 & P[e,x,x] \\
 & P[x,i[x],e] \\
 & P[i[x],x,e]
 \end{aligned}$$

We transform $\neg G$ into standard form

$$\begin{aligned}
& \neg \left(\left(\forall_x P[x, x, e] \right) \Rightarrow \left(\forall_{u,v,w} (P[u, v, w] \Rightarrow P[v, u, w]) \right) \right) \\
\iff & \neg \left(\neg \left(\forall_x P[x, x, e] \right) \vee \left(\forall_{u,v,w} (\neg P[u, v, w] \vee P[v, u, w]) \right) \right) \\
\iff & \left(\forall_x P[x, x, e] \right) \wedge \left(\exists_{u,v,w} (P[u, v, w] \wedge \neg P[v, u, w]) \right) \\
\rightsquigarrow & \forall_x P[x, x, e] \wedge P[a, b, c] \wedge \neg P[b, a, c]
\end{aligned}$$

which gives the following clauses

$$\begin{aligned}
& P[x, x, e] \\
& P[a, b, c] \\
& \neg P[b, a, c]
\end{aligned}$$

◀

Example 2 (Most General Unifier) Find a most general unifier for

$$W = \{P[a, x, f[g[y]]], P[z, f[z], f[u]]\}$$

Solution. Let $\sigma_0 = \varepsilon$ and $W_0 = W$. Since W_0 is not a singleton, σ_0 is not a mgu of W .

$$D_0 = \{a, z\}.$$

Let $\sigma_1 = \varepsilon \circ \{z \rightarrow a\}$, $W_1 = W_0\sigma_1 = \{P[a, x, f[g[y]]], P[a, f[a], f[u]]\}$.

W_1 is not a singleton. $D_1 = \{x, f[a]\}$.

Let $\sigma_2 = \{z \rightarrow a\} \{x \rightarrow f[a]\} = \{z \rightarrow a, x \rightarrow f[a]\}$. $W_2 = W_1\sigma_2 = \{P[a, f[a], f[g[y]]], P[a, f[a], f[u]]\}$.

W_2 is not a singleton. $D_2 = \{g[y], u\}$.

Let $\sigma_3 = \sigma_2 \{u \rightarrow g[y]\} = \{z \rightarrow a, x \rightarrow f[a], u \rightarrow g[y]\}$.

$W_3 = W_2\sigma_2 = \{P[a, f[a], f[g[y]]], P[a, f[a], f[g[y]]]\} = \{P[a, f[a], f[g[y]]]\}$.

Since W_3 is a singleton. $\sigma_3 = \{z \rightarrow a, x \rightarrow f[a], u \rightarrow g[y]\}$ is a mgu for W . ◀

Example 3 (Most General Unifier) Find a most general unifier for

$$W = \{Q[a], Q[b]\}$$

Solution. Let $\sigma_0 = \varepsilon$ and $W_0 = W$. Since W_0 is not a singleton, σ_0 is not a mgu of W .

$D_0 = \{a, b\}$. Since none of the elements of D_0 is a variable we conclude that W is not unifiable. ◀

Example 4 (Resolution 1) Prove by resolution the following

$$\forall_x F[x] \vee \forall_x H[x] \not\equiv \forall_x (F[x] \vee H[x])$$

Solution. Direction “ \Rightarrow ”. Let

$$\begin{aligned} F &: \iff \forall_x F[x] \vee \forall_x H[x] \\ G &: \iff \forall_x (F[x] \vee H[x]) \end{aligned}$$

We prove that G is a logical consequence of F by resolution. We have

$$\begin{aligned} F &: \iff \forall_x F[x] \vee \forall_x H[x] \\ &\iff \forall_{x,y} F[x] \vee H[y] \\ \neg G &: \iff \neg (\forall_x (F[x] \vee H[x])) \\ &\iff \exists_x (\neg F[x] \wedge \neg H[x]) \\ &\rightsquigarrow \neg F[a] \wedge \neg H[a] \end{aligned}$$

By transforming them into a set of clauses we have

$$\begin{aligned} (1) & F[x] \vee H[y] \\ (2) & \neg F[a] \\ (3) & \neg H[a] \end{aligned}$$

By applying resolution we obtain the following clauses

$$\begin{aligned} (4) & H[a] & (1) \wedge (2), \{x \rightarrow a, y \rightarrow a\} \\ (5) & \emptyset & (3) \wedge (4) \end{aligned}$$

Direction “ \Leftarrow ”. Let

$$\begin{aligned} F &: \iff \forall_x (F[x] \vee H[x]) \\ G &: \iff \forall_x F[x] \vee \forall_x H[x] \end{aligned}$$

We prove that G is a logical consequence of F by resolution. We have

$$\begin{aligned} F &: \iff \forall_x (F[x] \vee H[x]) \\ \neg G &: \iff \neg (\forall_x F[x] \vee \forall_x H[x]) \\ &\iff \exists_x \neg F[x] \wedge \exists_x \neg H[x] \\ &\rightsquigarrow \neg F[a] \wedge \neg H[b] \end{aligned}$$

By transforming them into a set of clauses we have

$$\begin{aligned} (1) & F[x] \vee H[x] \\ (2) & \neg F[a] \\ (3) & \neg H[b] \end{aligned}$$

By applying resolution we obtain the following clauses

$$\begin{array}{ll} (4) & H[a] \quad (1) \wedge (2), \{x \rightarrow a\} \\ (5) & F[b] \quad (1) \wedge (3), \{x \rightarrow b\} \end{array}$$

◀

Example 5 (Resolution 2) Prove by resolution that G is a logical consequence of F_1 and F_2 where

$$\begin{aligned} F_1 : & \forall_x (C[x] \Rightarrow (W[x] \wedge R[x])) \\ F_2 : & \exists_x (C[x] \wedge O[x]) \\ G : & \exists_x (O[x] \wedge R[x]) \end{aligned}$$

Solution. We show that $F_1 \wedge F_2 \wedge \neg G$ is unsatisfiable by resolution. We transform F_1 , F_2 , $\neg G$ into Skolem standard form. We have

$$\begin{aligned} F_1 : & \forall_x (C[x] \Rightarrow (W[x] \wedge R[x])) \\ \iff & \forall_x (\neg C[x] \vee (W[x] \wedge R[x])) \\ \iff & \forall_x (\neg C[x] \vee W[x]) \wedge (\neg C[x] \vee R[x]) \\ \\ F_2 : & \exists_x (C[x] \wedge O[x]) \\ \rightsquigarrow & C[a] \wedge O[a] \\ \\ \neg G : & \neg (\exists_x (O[x] \wedge R[x])) \\ \iff & \forall_x (\neg O[x] \vee \neg R[x]) \end{aligned}$$

We have the following set of clauses

$$\begin{array}{ll} (1) & \neg C[x] \vee W[x] \\ (2) & \neg C[x] \vee R[x] \\ (3) & C[a] \\ (4) & O[a] \\ (5) & \neg O[x] \vee \neg R[x] \end{array}$$

By resolution we obtain also the following clauses

$$\begin{array}{ll} (6) & \neg R[a] \quad (4) \wedge (5), \{x \rightarrow a\} \\ (7) & \neg C[a] \quad (6) \wedge (2), \{x \rightarrow a\} \\ (8) & \emptyset \quad (7) \wedge (3) \end{array}$$

◀

Example 6 (Resolution 3) Prove by resolution that G is a logical consequence of F_1 and F_2 where

$$\begin{aligned} F_1 &: \exists_x \left(P[x] \wedge \forall_y (D[y] \Rightarrow L[x, y]) \right) \\ F_2 &: \forall_x \left(P[x] \Rightarrow \forall_y (Q[y] \Rightarrow \neg L[x, y]) \right) \\ G &: \forall_x (D[x] \Rightarrow \neg Q[x]) \end{aligned}$$

Solution. We show that $F_1 \wedge F_2 \wedge \neg G$ is unsatisfiable by resolution. We transform F_1 , F_2 , $\neg G$ into Skolem standard form. We have

$$\begin{aligned} F_1 &: \exists_x \left(P[x] \wedge \forall_y (D[y] \Rightarrow L[x, y]) \right) \\ &\iff \exists_x \left(P[x] \wedge \forall_y (\neg D[y] \vee L[x, y]) \right) \\ &\iff \exists_{xy} (P[x] \wedge (\neg D[y] \vee L[x, y])) \\ &\rightsquigarrow \forall_y (P[a] \wedge (\neg D[y] \vee L[a, y])) \\ \\ F_2 &: \forall_x \left(P[x] \Rightarrow \forall_y (Q[y] \Rightarrow \neg L[x, y]) \right) \\ &\iff \forall_x \left(P[x] \Rightarrow \forall_y (\neg Q[y] \vee \neg L[x, y]) \right) \\ &\iff \forall_x \left(\neg P[x] \vee \forall_y (\neg Q[y] \vee \neg L[x, y]) \right) \\ &\iff \forall_{xy} (\neg P[x] \vee \neg Q[y] \vee \neg L[x, y]) \\ \\ \neg G &: \neg \left(\forall_x (D[x] \Rightarrow \neg Q[x]) \right) \\ &\iff \neg \left(\forall_x (\neg D[x] \vee \neg Q[x]) \right) \\ &\iff \exists_x (D[x] \wedge Q[x]) \\ &\rightsquigarrow D[a] \wedge Q[a] \end{aligned}$$

We have the following set of clauses

- (1) $P[a]$
- (2) $\neg D[y] \vee L[a, y]$
- (3) $\neg P[x] \vee \neg Q[y] \vee \neg L[x, y]$
- (4) $D[a]$
- (5) $Q[a]$

By resolution we obtain also the following clauses

$$\begin{array}{ll}
 (6) & L[a, a] \\
 (7) & \neg P[a] \vee \neg Q[a] \\
 (8) & \neg Q[a] \\
 (9) & \emptyset
 \end{array}
 \quad
 \begin{array}{ll}
 (2) \wedge (4), \{y \rightarrow a\} \\
 (3) \wedge (6), \{x \rightarrow a, y \rightarrow a\} \\
 (1) \wedge (7) \\
 (5) \wedge (8)
 \end{array}$$

◀

Example 7 (Resolution 4) Prove by resolution that G is a logical consequence of F where

$$\begin{aligned}
 F : & \underset{x,y}{\forall \exists} (S[x, y] \wedge M[y]) \Rightarrow \underset{y}{\exists} (I[y] \wedge E[x, y]) \\
 G : & \underset{x}{\neg \exists} I[x] \Rightarrow \underset{x,y}{\forall} (S[x, y] \Rightarrow \neg M[y])
 \end{aligned}$$

Solution. We show that $F \wedge \neg G$ is unsatisfiable. First we transform the formulae into standard form. We have

$$\begin{aligned}
 F : & \underset{x}{\forall} \left(\underset{y}{\exists} (S[x, y] \wedge M[y]) \right) \Rightarrow \underset{y}{\exists} (I[y] \wedge E[x, y]) \\
 \iff & \underset{x}{\forall} \neg \left(\underset{y}{\exists} (S[x, y] \wedge M[y]) \right) \vee \underset{y}{\exists} (I[y] \wedge E[x, y]) \\
 \iff & \underset{x}{\forall} \left(\underset{y}{\forall} (\neg S[x, y] \vee \neg M[y]) \right) \vee \underset{y}{\exists} (I[y] \wedge E[x, y]) \\
 \iff & \underset{x}{\forall} \left(\underset{y}{\forall} (\neg S[x, y] \vee \neg M[y]) \right) \vee (I[f[x]] \wedge E[x, f[x]]) \\
 \iff & \underset{x,y}{\forall \forall} (\neg S[x, y] \vee \neg M[y]) \vee (I[f[x]] \wedge E[x, f[x]]) \\
 \iff & \underset{x,y}{\forall \forall} ((\neg S[x, y] \vee \neg M[y] \vee I[f[x]]) \wedge (\neg S[x, y] \vee \neg M[y] \vee E[x, f[x]])) \\
 \neg G : & \neg \left(\underset{x}{\neg \exists} I[x] \Rightarrow \underset{x,y}{\forall} (S[x, y] \Rightarrow \neg M[y]) \right) \\
 \iff & \neg \left(\underset{x}{\neg \exists} I[x] \Rightarrow \underset{x,y}{\forall} (\neg S[x, y] \vee \neg M[y]) \right) \\
 \iff & \neg \left(\underset{x}{\exists} I[x] \vee \underset{x,y}{\forall} (\neg S[x, y] \vee \neg M[y]) \right) \\
 \iff & \left(\underset{x}{\forall} \neg I[x] \wedge \underset{x,y}{\exists} (S[x, y] \wedge M[y]) \right) \\
 \iff & \underset{z}{\forall} \neg I[z] \wedge \underset{x,y}{\exists} (S[x, y] \wedge M[y]) \\
 \rightsquigarrow & \underset{z}{\forall} \neg I[z] \wedge S[a, b] \wedge M[b]
 \end{aligned}$$

We have the following set of clauses

- (1) $\neg S[x, y] \vee \neg M[y] \vee I[f[x]]$
- (2) $\neg S[x, y] \vee \neg M[y] \vee E[x, f[x]]$
- (3) $\neg I[z]$
- (4) $S[a, b]$
- (5) $M[b]$

By resolution we obtain also the following clauses

- (6) $\neg S[x, y] \vee \neg M[y] \quad (1) \wedge (3), \{z \rightarrow f[x]\}$
- (7) $\neg M[b] \quad (4) \wedge (6), \{x \rightarrow a, y \rightarrow b\}$
- (8) $\emptyset \quad (5) \wedge (7)$

◀

Example 8 (Resolution 5) Prove by resolution that G is a logical consequence of F_1, F_2 , and F_3 where

$$\begin{aligned} F_1 : & \forall_x (Q[x] \Rightarrow \neg P[x]) \\ F_2 : & \forall_x ((R[x] \wedge \neg Q[x]) \Rightarrow \exists_y (T[x, y] \wedge S[y])) \\ F_3 : & \exists_x (P[x] \wedge \forall_y (T[x, y] \Rightarrow P[y]) \wedge R[x]) \\ G : & \exists_x (S[x] \wedge P[x]) \end{aligned}$$

Solution. We show that $F_1 \wedge F_2 \wedge F_3 \wedge \neg G$ is unsatisfiable. First we transform the formulae into standard form.

$$\begin{aligned}
F_1 : \quad & \forall_x (Q[x] \Rightarrow \neg P[x]) \iff \forall_x (\neg Q[x] \vee \neg P[x]) \\
F_2 : \quad & \forall_x \left((R[x] \wedge \neg Q[x]) \Rightarrow \exists_y (T[x, y] \wedge S[y]) \right) \\
& \iff \forall_x \left(\neg(R[x] \wedge \neg Q[x]) \vee \exists_y (T[x, y] \wedge S[y]) \right) \\
& \iff \forall_x \left(\neg R[x] \vee Q[x] \vee \exists_y (T[x, y] \wedge S[y]) \right) \\
& \iff \forall_{xy} (\neg R[x] \vee Q[x] \vee (T[x, y] \wedge S[y])) \\
& \iff \forall_{xy} ((\neg R[x] \vee Q[x] \vee T[x, y]) \wedge (\neg R[x] \vee Q[x] \vee S[y])) \\
& \rightarrow \forall_x ((\neg R[x] \vee Q[x] \vee T[x, f[x]]) \wedge (\neg R[x] \vee Q[x] \vee S[f[x]])) \\
F_3 : \quad & \exists_x \left(P[x] \wedge \forall_y (T[x, y] \Rightarrow P[y]) \wedge R[x] \right) \\
& \iff \exists_x \left(P[x] \wedge \forall_y (\neg T[x, y] \vee P[y]) \wedge R[x] \right) \\
& \iff \exists_{xy} (P[x] \wedge (\neg T[x, y] \vee P[y]) \wedge R[x]) \\
& \rightarrow \forall_y (P[a] \wedge (\neg T[a, y] \vee P[y]) \wedge R[a]) \\
\neg G : \neg & \left(\exists_x (S[x] \wedge P[x]) \right) \\
& \iff \forall_x (\neg S[x] \vee \neg P[x])
\end{aligned}$$

From the normal forms we obtain the clauses (1) to (7), and then by resolution we obtain the clauses (8) to (15):

(1)	$\neg Q[x] \vee \neg P[x]$	F_1
(2)	$\neg R[x] \vee Q[x] \vee T[x, f[x]]$	F_2
(3)	$\neg R[x] \vee Q[x] \vee S[f[x]]$	F_2
(4)	$P[a]$	F_3
(5)	$\neg T[a, y] \vee P[y]$	F_3
(6)	$R[a]$	F_3
(7)	$\neg S[x] \vee \neg P[x]$	$\neg G$
(8)	$\neg Q[a]$	(1) \wedge (4), $\{x \rightarrow a\}$
(9)	$\neg R[a] \vee T[a, f[a]]$	(8) \wedge (2), $\{x \rightarrow a\}$
(10)	$\neg R[a] \vee P[f[a]]$	(9) \wedge (5), $\{y \rightarrow f[a]\}$
(11)	$P[f[a]]$	(10) \wedge (6)
(12)	$\neg S[f[a]]$	(11) \wedge (7)
(13)	$\neg R[a] \vee Q[a]$	(12) \wedge (3)
(14)	$Q[a]$	(13) \wedge (6)
(15)	\emptyset	(14) \wedge (8)

◀